

Generalized second law of thermodynamics in the emergent universe for some viable models of $f(T)$ gravity

Rahul Ghosh^{1,*} and Surajit Chattopadhyay^{2,†}

¹*Department of Mathematics, Bhairab Ganguly College, Kolkata-700 056, India.*

²*Pailan College of Management and Technology,
Bengal Pailan Park, Kolkata-700 104, India.*

(Dated: July 26, 2012)

The present study is motivated by the study of reference [1], where the generalized second law of thermodynamics has been investigated for a flat FRW universe for two viable models of $f(T)$ gravity. In the present work, we have considered a non-flat universe and accordingly studied the behaviors of equation of state parameter and deceleration parameter. Subsequently, using the first law of thermodynamics we derived the expressions for the time derivative of the total entropy of a universe enveloped by apparent horizon. In the next phase, with the choice of scale factor pertaining to an emergent universe we have investigated the sign of the time derivatives of total entropy for three viable models of $f(T)$ gravity.

I. INTRODUCTION

The cosmic acceleration today has been supported by independent observational data and the origin of dark energy responsible for this cosmic acceleration is one of the most serious problems in modern cosmology the modification of gravity on large scales [2]. Motivated by attempts to explain the observed acceleration of the universe in a natural way, there has been a great deal of recent interest in a generalization of this theory in which the Lagrangian is an arbitrary algebraic function f of the Lagrangian of teleparallel gravity T [3]. This is similar to creating $f(R)$ gravity theories that are a generalization of general relativity (review on $f(R)$ gravity is available in [4]). This gravity is dubbed as $f(T)$ gravity. Review on $f(T)$ gravity is available in the references [5–8]. This gravity describes the present accelerating expansion of the universe without resorting to dark energy. It is a generalization of the teleparallel gravity (TG) by replacing the so-called torsion scalar T with $f(T)$ [1]. The possibilities of an emergent universe [9–11] have been studied recently in a number of papers in which one looks for a universe which is ever-existing and large enough

*Electronic address: ghoshrahul3@gmail.com

†Electronic address: surajit.2008@yahoo.co.in, surajcha@iucaa.ernet.in

so that the spacetime may be treated as classical entities. In these models, the universe in the infinite past is in an almost static state but it eventually evolves into an inflationary stage [11]. An emergent universe model can be defined as a singularity free universe which is ever existing with an almost static nature in the infinite past ($t \rightarrow -\infty$) and then evolves into an inflationary stage [12]. In the references [11, 13, 14] the characteristics of an emergent universe have been summarized as:

1. the universe is almost static at the finite past ($t \rightarrow -\infty$) and isotropic, homogeneous at large scales;
2. it is ever existing and there is no timelike singularity;
3. the universe is always large enough so that the classical description of space-time is adequate;
4. the universe may contain exotic matter so that the energy conditions may be violated;
5. the universe is accelerating as suggested by recent measurements of distances of high redshift type Ia supernovae.

In the reference [14], the equation of state parameter was reconstructed for emergent universe under $f(T)$ gravity. In another work, reference [13] considered the generalized Ricci dark energy and generalized holographic dark energy in the scenario of an emergent universe and studied the behaviors of the potential and the chameleon scalar field. In the references [12–14], the choice of scale factor a for emergent universe was $a = a_0 (\eta + e^{Bt})^n$, where $a_0 > 0$, $\eta > 0$, $B > 0$ and $n > 0$.

The present work is motivated by the work of [1], who investigated the validity of the generalized second law (GSL) of gravitational thermodynamics in the framework of $f(T)$ gravity. The present study deviates the earlier work in the following aspects: (i) we have considered three viable models of $f(T)$ gravity without the assumption of flat universe and examined the cases of non-flat universe; (ii) we have examined the validity of GSL of thermodynamics for a particular choice of scale factor pertaining to emergent universe; (iii) we have investigated some particular cases for the violation of GSL.

II. $f(T)$ GRAVITY IN NON-FLAT UNIVERSE

In the framework of $f(T)$ theory, the action of modified TG is given by [1]

$$I = \frac{1}{16\pi G} \int d^4x \sqrt{-g} [f(T) + L_m] \quad (1)$$

where, L_m is the Lagrangian density of the matter inside the universe. We consider a Friedmann-Robertson-Walker (FRW) universe filled with the pressureless matter. Choosing ($8\pi G = 1$) the modified Friedman equations in the framework of $f(T)$ gravity are given by [1, 15]

$$H^2 + \frac{k}{a^2} = \frac{1}{3}(\rho_m + \rho_T) \quad (2)$$

$$\dot{H} - \frac{k}{a^2} = -\frac{1}{2}(\rho_m + \rho_T + p_T) \quad (3)$$

where,

$$\rho_T = \frac{1}{2}(2Tf_T - f - T) \quad (4)$$

$$p_T = -\frac{1}{2} \left[-8\dot{H}Tf_{TT} + (2T - 4\dot{H})f_T - f + 4\dot{H} - T \right] \quad (5)$$

Where [15]

$$T = -6 \left(H^2 + \frac{k}{a^2} \right) \quad (6)$$

Here, k is the curvature of the universe and $H = \frac{\dot{a}}{a}$ is the Hubble parameter and ρ_m is the energy density of the matter. Also ρ_T and p_T are the torsion contributions to the energy density and pressure. Energy conservation equations are given by

$$\dot{\rho}_m + 3H(\rho_m) = 0; \quad \dot{\rho}_T + 3H(\rho_T + p_T) = 0 \quad (7)$$

The effective equation of state (EoS) parameter due to the torsion contribution is defined as [1]

$$w_T = -1 - \frac{\dot{T}}{3H} \left(\frac{2Tf_{TT} + f_T - 1}{2Tf_T - f - T} \right) \quad (8)$$

From equations (2),(4)and (6) we get ρ_m as

$$\rho_m = \frac{1}{2}(f - 2Tf_T) \quad (9)$$

Using (3) and (9) we get the time derivative of Hubble parameter as

$$\dot{H} = -\frac{1}{2} \left(\frac{\rho_m - \frac{2k}{a^2}}{f_T + 2Tf_{TT}} \right) \quad (10)$$

Using equations (6) and (10) we get

$$\dot{T} = \frac{12H}{f_T + 2Tf_{TT}} \left[\frac{k(f_T + 2Tf_{TT} - 1)}{a^2} + \frac{\rho_m}{2} \right] \quad (11)$$

Using (2), (9) and (10) we get the deceleration parameter as

$$q = -1 - \frac{\dot{H}}{H^2} = -1 + \frac{3}{2} \left[\frac{\rho_m a^2 - 2k}{(f_T + 2Tf_{TT})(a^2(\rho_m + \rho_T) - 3k)} \right] \quad (12)$$

III. GSL IN $f(T)$ GRAVITY

The apparent horizon has been argued as a causal horizon for a dynamical spacetime and is associated with gravitational entropy and surface gravity [16]. The radius of the apparent horizon, which is denoted by R_A , is given by [17]

$$R_A = \frac{1}{\sqrt{H^2 + \frac{k}{a^2}}} \quad (13)$$

In the present work we are investigating the validity of GSL in a non-flat FRW universe filled with pressureless dark matter. The GSL can be expressed as $\dot{S}_A + \dot{S}_m \geq 0$, where S_A denotes the Bekenstein-Hawking entropy on the horizon and S_m is the entropy due to the matter sources inside the horizon [18]. Detailed account of the GSL is available in [17, 19, 20]. From the first law of thermodynamics one can get (Clausius relation) $-dE = T_A dS_A$ to the apparent horizon R_A . The Friedmann equation in the Einstein gravity can be derived if we take the Hawking temperature $T_A = 1/2\pi R_A$ and the entropy $S_A = A/4G$ on the apparent horizon, where A is the area of the horizon [1]. However, this definition is changed for other modified gravity theories. In $f(T)$ gravity, it was shown that when f_{TT} is small, the first law of black hole thermodynamics is satisfied approximatively and the entropy of horizon is $S_A = \frac{A f_T}{4G}$, where f_T denotes the derivative with respect to the torsion scalar T [1]. In the present work we assume that the boundary of the universe to be enclosed by the dynamical apparent horizon R_A and the Hawking temperature on the apparent horizon is given by [1, 17]

$$T_A = \frac{1}{2\pi R_A} \left(1 - \frac{\dot{R}_A}{2H R_A} \right) \quad (14)$$

Based on the discussions in the last section we compute

$$T_A \dot{S}_m = 2\pi R_A^2 (f - 2T f_T) (\dot{R}_A - H R_A) \quad (15)$$

$$T_A \dot{S}_A = 4\pi \left(1 - \frac{\dot{R}_A}{2H R_A} \right) \left[2\dot{R}_A f_T + 6H \left(\frac{\rho_m - \frac{2k}{a^2}}{f_T + 2T f_{TT}} \right) R_A f_{TT} \right] \quad (16)$$

where, $\dot{S}_m$ and $\dot{S}_A$ denote the time derivatives of the entropy due to the matter sources inside the horizon and on the horizon respectively. In the next subsection we investigate whether $\dot{S}_m + \dot{S}_A \geq 0$ for three viable choices of $f(T)$ with the choice of scale factor for emergent scenario, i.e. $a = a_0 (\eta + e^{Bt})^n$, where $a_0 > 0$, $\eta > 0$, $B > 0$ and $n > 0$. The $f(T)$ models considered here are

1. $f(T) = \gamma T + \lambda T^m$ [14]

2. $f(T) = T - \lambda T \left(1 - e^{\left(\beta \frac{T_0}{T}\right)}\right)$ where $\lambda = \frac{1-8\pi \frac{\rho_{m0}}{3H_0^2}}{1-(1-2\beta)e^\beta} [1]$
3. $f(T) = T + \lambda(-T)^{n_1}$ where $\lambda = \left(\frac{1-8\pi \frac{\rho_{m0}}{3H_0^2}}{2n_1-1}\right) (6H_0^2)^{(1-n_1)} [1]$

A. Model I

First we consider $f(T) = \gamma T + \lambda T^m$. In this case we have

$$\dot{T} = \frac{\xi_1}{\xi_2} \quad (17)$$

where,

$$\begin{aligned} \xi_1 = & 6Be^{Bt}n(e^{Bt} + \eta)^{-3-2n} [6(k^2(2 + \gamma)(e^{Bt} + \eta)^4 + 3a_0^4B^4e^{4Bt}n^4\gamma \times \\ & (e^{Bt} + \eta)^{4n} - 2a_0^2B^2e^{2Bt}kn^2(1 + 2\gamma)(e^{Bt} + \eta)^{2+2n}) + a_0^2(e^{Bt} + \eta)^{2n} \times \\ & \left\{ -\frac{B^2e^{2Bt}n^2}{(e^{Bt} + \eta)^2} + \frac{k(e^{Bt} + \eta)^{-2n}}{a_0^2} \right\}^m (-6^mk(3 + 4(-2 + m)m)(e^{Bt} + \eta)^4 + 3a_0^2B^2e^{2Bt})\lambda] \end{aligned} \quad (18)$$

$$\begin{aligned} \xi_2 = & a_0^2 [-6k\gamma(e^{Bt} + \eta)^2 + 6a_0^2B^2e^{2Bt}n^2\gamma(e^{Bt} + \eta)^{2n} + a_0^2m(6^m - 2^{1+m}3^m m)(e^{Bt} + \eta)^{2+2n}\lambda \times \\ & \left\{ -\frac{B^2e^{2Bt}n^2}{(e^{Bt} + \eta)^2} + \frac{k(e^{Bt} + \eta)^{-2n}}{a_0^2} \right\}^m] \end{aligned} \quad (19)$$




FIG. 1: Evolution of $\dot{T}$ with cosmic time t for model I.







FIG. 2: Behavior of w_T for model I.

In figure 1 we have viewed the behavior of the time derivative of T for $f(T) = \gamma T + \lambda T^m$ and it is found to be an increasing function of time. The EoS parameter for this model has been plotted in figure 2, where it is observed that the EoS parameter $w_T \leq -1$ that indicates phantom like behavior. Also it is observed that from early to late stage of the universe the EoS parameter is tending towards -1 . This holds for $k = -1, +1$ and 0 . Deceleration parameter q plotted in figure 3 remains negative throughout the evolution of the universe for model I. This indicates ever-accelerating universe. For

FIG. 3: Behavior of $p_T + \rho_T$ for model I.FIG. 4: Behavior of q_T for model I.FIG. 5: Behavior of $\dot{S}_{total}$ when the apparent horizon is considered as the enveloping horizon under $f(T) = \gamma T + \lambda T^m$ i.e. model I.

all of the above plots we have taken $B = 2$, $a_0 = 1.5$, $n = 1.2$, $\eta = 3$, $\gamma = 4.02$, $\lambda = 4.03$, and $m = 2$. The sum of energy density and pressure is placed in figure 4, where the sum is found to be negative. This indicates the violation of the strong energy condition. In figure 5, we have plotted the time derivative of total entropy $\dot{S}_{total}$ against cosmic time t . The $\dot{S}_{total}$ is computed using equations (15) and (16), where $f(T)$ has been replaced by the form of model I. The $\dot{S}$ is found to be positive throughout the evolution of the universe. This indicates that the GSL is satisfied by model I under emergent scenario of the universe.

B. Model II

In this section we consider model II where $f(T) = T - \lambda T \left(1 - e^{\left(\beta \frac{T_0}{T} \right)} \right)$ where $\lambda = \frac{1 - 8\pi \frac{\rho_{m0}}{3H_0^2}}{1 - (1 - 2\beta)e^\beta}$. In this case first we view the behavior of the torsion scalar with the evolution of the universe. The time derivative of T is plotted against cosmic time t in figure 1 and a behavior similar to that for model I is observed. It may be noted that for model II our choices are $a_0 = 20$, $B = 4$, $n = 0.04$ and

$\eta = 3$ that satisfy the conditions for emergent universe. For the form of $f(T)$ under consideration, the choices are $\lambda = 0.03$, $\beta = -0.02$, $\rho_{m0} = 0.23$ and $H_0 = 74.2$. In all the figures, the red, green and blue lines correspond to the cases $k = -1$, $k = +1$ and $k = 0$ respectively. Like model I,




FIG. 6: Behavior of $\dot{T}$ for model II.




FIG. 7: Behavior of w_T for model II.

the $\dot{T}$ (figure 6) stays positive and exhibits upward movement with evolution of the universe. The EOS parameter, as seen in figure 7, stays below -1 and does never tend to -1 . This indicates phantom-like behavior of the EOS parameter. The deceleration parameter remains negative for every choice of k . Figure 8, where $(p_T + \rho_T)$ is plotted against t shows the violation of strong energy




FIG. 8: Behavior of $(p_T + \rho_T)$ for model II.




FIG. 9: Behavior of q_T for model II.

condition by model II. The accelerated universe under model II is understandable from figure 9 that shows negative q_T . Finally, when we consider the time derivative of total entropy $\dot{S}$ in figure 10, we find that it remains negative for every choice of k . This indicates violation of GSL by model II.




FIG. 10: Behavior of $\dot{S}_{total}$ when the apparent horizon is considered as the enveloping horizon under $f(T) = T - \lambda T \left(1 - e^{\left(\beta \frac{T_0}{T}\right)}\right)$ i.e. model II.




FIG. 11: Behavior of $\dot{T}$ for model III i.e. $f(T) = T + \lambda(-T)^{n_1}$. The solid lines pertain to $n_1 < 1$ and the dashed lines pertain to $n_1 > 1$.

C. Model III

In this section we consider model III where $f(T) = T + \lambda(-T)^{n_1}$ where $\lambda = \left(\frac{1-8\pi\frac{\rho_{m0}}{3H_0^2}}{2n_1-1}\right)(6H_0^2)^{(1-n_1)}$. To study the model III, we have taken both $n_1 < 1$ ($n_1 = \frac{1}{3}$) and $n_1 > 1$ ($n_1 = 1.2$) into account. Accordingly we have plotted $\dot{T}$ in figure 11, where the $\dot{T}$ is found to behave in the manner similar to the cases of models I and II. From figures 12 and 13 we find that the EOS parameter w_T behave like phantom and tend to -1 in both cases irrespective of the choice of k . The deceleration parameters plotted in figures 14 and 15 indicated the realization of the accelerated expansion of the universe by model III for $n_1 < 1$ and $n_1 > 1$ respectively.

Finally when we plot $\dot{S}_{total}$ in figures 16 and 17 for the case of model III with apparent horizon as the enveloping horizon, we observe violation of the GSL by model III irrespective of the choices of n_1 .




FIG. 12: Behavior of w_T for model III with $n_1 < 1$.




FIG. 13: Behavior of w_T for model III with $n_1 > 1$.




FIG. 14: Behavior of q_T for model III with $n_1 < 1$.




FIG. 15: Behavior of q_T for model III with $n_1 > 1$.

IV. DISCUSSIONS

In this work we have considered three viable $f(T)$ gravity with scale factor a pertaining to emergent universe. We have investigated the validity of GSL for all the cases along with some related characteristics. It is observed that for model I the time derivative of T exhibits an increasing pattern in the positive side with evolution of the universe. When we considered the EOS parameter for this model we find that with evolution of the universe the w_T remains below -1 at earlier stages and tends to -1 at later stage. This indicates phantom-like behavior of w_T . This is displayed in figure 2. In figure 3 we have plotted $p_T + /rho_T$ against cosmic time t and we find that it is staying at negative level. This indicates violation of strong energy condition. The deceleration parameter q_T for this model is plotted against cosmic time t in figure 4 and we find that it is negative throughout. This indicates accelerated phase of the universe. The time derivative of total entropy is plotted in figure 5 and we find that it remains positive through the evolution of the universe irrespective of $k = -1, +1$ or 0 . Therefore our observations are that for model I are: (i) the EOS exhibits phantom-like behavior, (ii) The strong energy condition is violated, (iii) we get




FIG. 16: Behavior of $\dot{S}_{total}$ when the apparent horizon is considered as the enveloping horizon under $f(T) = T + \lambda(-T)^{n_1}$ with $n_1 < 1$.




FIG. 17: Behavior of $\dot{S}_{total}$ when the apparent horizon is considered as the enveloping horizon under $f(T) = T + \lambda(-T)^{n_1}$ with $n_1 > 1$.

an accelerating universe and (iv) generalized second law of thermodynamics holds irrespective of the curvature of the universe. It should be stated that the apparent horizon has been considered as the enveloping horizon of the universe. Now we consider the model II with the choices of scale factor and enveloping horizon similar to that of model I. All the quantities considered for model I are considered for model II also. From figure 6 we observe that the behavior of the time derivative of T is almost similar to that of model I. However the EOS parameter, as plotted in 7, exhibits a little difference with model I. In the present case the EOS parameter exhibits phantom-like behavior. However it does tend to -1 at later stages of the universe. The strong energy condition is violated like model I (figure 8). Like model I here also we find an ever accelerating universe (see figure 9). Finally when we consider the GSL we find a prominent difference between model I and model II. In model II the time derivative of total entropy remains negative through the evolution of the universe. This indicates the violation the GSL. This is displayed in figure 10. Therefore for model II the observations are following: (i) the EOS exhibits phantom-like behavior, (ii) The strong energy condition is violated, (iii) we get accelerating universe and (iv) generalized second law of thermodynamics is violated here. Here also the above results hold irrespective of the choice of k . Now we come to Model III. For model III we have same choices of scale factor and enveloping horizon. However, while considering model III we consider two cases, namely $n_1 < 1$ and $n_1 > 1$. From figure 11 we find that the time derivative of T remains positive for $n_1 < 1$ as well as $n_1 > 1$ and in both cases it shows increasing pattern. In figure 12 we have plotted EOS parameter for $n_1 < 1$ and we have observed that it is always staying below -1 irrespective of the choice of curvature. This indicates phantom-like behavior of the EOS parameter for $n_1 < 1$. In figure 13, we plot the EOS for $n_1 > 1$ and here also we observe phantom like behavior. In

figure 14 and 15 we plot the deceleration parameter q_T for $n_1 < 1$ and $n_1 > 1$ respectively and in both of the cases we get negative deceleration parameter indicating ever accelerating universe. Now we come to the GSL. The GSL for $n_1 < 1$ is considered in figure 16 and we observe that the time derivative of total entropy is staying at negative level. This means the violation of the GSL. The $n_1 > 1$ case is considered in figure 17. Here also we find that the time derivative of the total entropy is negative through the universe. Therefore we conclude that the GSL is not valid in this case. The above results holds for $k = -1, +1$ and 0 . Therefore our observations are the following: (i) the EOS behaves like phantom, (ii) accelerated expansion of the universe is realized and (iii) the generalized second law of thermodynamics does not work here. The above results are independent of the choice of k .

V. ACKNOWLEDGEMENTS

The first author wishes to thank the Inter-University Centre for Astronomy and Astrophysics (IUCAA), Pune, India for providing warm hospitality during a scientific visit in January 2012, when part of the work was carried out. The second author sincerely acknowledges the Visiting Associateship provided by IUCAA, Pune, India for the period of August 2011 to July 2014 to carry out research in General Relativity and Cosmology.

-
- [1] K. Karami and A. Abdolmaleki, JCAP **04** (2012) 007.
 - [2] S. Tsujikawa, Lect. Notes Phys. **800**(2010) 99.
 - [3] B. Li, T. P. Sotiriou and J. D. Barrow, Phys. Rev. D **83** (2011) 064035.
 - [4] T. P. Sotiriou and V. Faraoni, Rev. Mod. Phys. **82** (2010) 451.
 - [5] R. Zheng and Q-G. Huang, JCAP **03** (2011) 002.
 - [6] J. B. Dent, S. Dutta and E. N. Saridakis, JCAP **01** (2011) 009.
 - [7] G. R. Bengochea, Phys. Lett. B **695** (2011) 405.
 - [8] R. Myrzakulov, Eur. Phys. J. C **71** (2011) 1752.
 - [9] G. F. R. Ellis and R. Maartens, Class. Quantum Grav. **21** (2004) 223.
 - [10] G. F. R. Ellis, J. Murugan and C. G. Tsagas, Class. Quantum Grav. **21** (2004) 233.

- [11] S. Mukherjee, B. C. Paul, N. K. Dadhich, S. D. Maharaj and A. Beesham, *Class. Quantum Grav.* **23** (2006) 6927.
- [12] U. Debnath, *Class. Quantum Grav.* **25** (2008) 205019.
- [13] S. Chattopadhyay and U. Debnath, *Can. J. Phys.* **89** (2011) 941.
- [14] S. Chattopadhyay and U. Debnath, *Int. J. Mod. Phys. D* **20** (2011) 1135.
- [15] R. Ferraro and F. Fiorini, **702** (2011) *Phys. Lett. B* 75.
- [16] D. Bak and S. J. Rey, **17** (2000) *Class. Quantum Grav.* L83.
- [17] A. Sheykhi, **27** (2010) *Class. Quantum Grav.* 025007.
- [18] M. Akbar, **25** *Chin. Phys. Lett.* 4199.
- [19] M. Jamil, E. N. Saridakis and M. R. Setare, **11** (2010) *JCAP* 032.
- [20] T. Padmanabhan, **73** (2010) 046901